



其中地中海生境取自 7 个种，萨瓦纳生境取自 2 个种，热带雨林生境取自 8 个种，红树植物数据来自本研究。

Among them, 7 species were collected from Mediterranean habitat, 2 species from Savannah habitat, 8 species from tropical rainforest habitat, and mangrove plant data were obtained from this study.

附图 1 热带、亚热带不同地区常绿植物叶黄素库容量

Fig. 1 Xanthophyll pool size of evergreen plants in tropical and subtropical regions

参考文献:

- [1]Faria T, Silverio D, Breia E, Cabral R, Abadia A, *et al.* Differences in the response of carbon assimilation to summer stress (water deficits, high light and temperature) in four Mediterranean tree species[J]. *Physiol Plantarum*, 1998, 102(3): 419-428
- [2]Zhu JJ, Zhang JL, Liu HC, Cao KF. Photosynthesis, non - photochemical pathways and activities of antioxidant enzymes in a resilient evergreen oak under different climatic conditions from a valley - savanna in Southwest China[J]. *Physiol Plantarum*, 2009, 135(1): 62-72
- [3]García-Plazaola JI, Esteban R, Hormaetxe K, Fernández-Marín B, Becerril JM. Photoprotective responses of Mediterranean and Atlantic trees to the extreme heat-wave of summer 2003 in Southwestern Europe[J]. *Trees*, 2008, 22(3): 385-392
- [4]Franco AC, Matsubara S, Orthen B. Photoinhibition, carotenoid composition and the co-regulation of photochemical and non-photochemical quenching in neotropical savanna trees[J]. *Tree Physiol*, 2007, 27(5): 717-725
- [5]Matsubara S, Krause GH, Aranda J, Virgo A, Beisel KG, Jahns P, Winter K. Sun-shade patterns of leaf carotenoid composition in 86 species of neotropical forest plants[J]. *Funct Plant Biol*, 2007, 36(1): 20-36